



Working Women Behavior On Cervical Cancer As Participants Of National Health Insurance

Perilaku Wanita Pekerja Terhadap Kanker Serviks Sebagai Peserta Jaminan Kesehatan Nasional

Awang Saputra¹, Suginarti², Ahmad Faridi³

¹ Health Service Office of Serang City, Banten, Indonesia/Public Health Magister Program University of Muhammadiyah Prof.Dr.HAMKA, Jakarta, Indonesia

² Public Health Postgraduate Program University of Muhammadiyah Prof.Dr.HAMKA, Jakarta, Indonesia

³ Nutrition Science Department University of Muhammadiyah Prof.Dr.HAMKA, Jakarta, Indonesia

ABSTRACT

One indicator to determine the high degree of health of a country showed from maternal mortality (MMR). For some areas that have difficult access to health services, the Ministry of Health makes a policy by developing partnership programs for health workers and traditional birth attendants. The phenomenon of traditional birth attendants becomes an interesting phenomenon because of the values of local wisdom they guard. This study aims to find out more about the experience of participants at-home birth with a partnership of health workers and traditional birth attendants. The method used with qualitative design with phenomenology approach. Participants were determined using Purposive sampling method. Methods of data collection used in-depth interviews with mothers and FGDs on Kepala Desa, Pemangku Adat, traditional birth attendants, and health workers. The results of this study obtained 6 themes of choice of place and helpers during childbirth, mother beliefs against traditional birth attendants in the process of childbirth, Tradition of manyirira and sequential during pregnancy and after childbirth, mother's feelings with partnership, independent task of traditional birth attendants after childbirth and Utilization of health services. Suggestion in this research is health workers must know culture adopted by patient and family cared. It aims to give an overview and positive impact on the enhancement of the quality of maternal health services in the context of culture.

ABSTRAK

Salah satu indikator untuk menentukan tingginya derajat kesehatan suatu negara dilihat dari angka kematian ibu (AKI). Untuk beberapa daerah yang memiliki akses pelayanan kesehatan yang sulit, Kementerian Kesehatan membuat sebuah kebijakan dengan mengembangkan program kemitraan tenaga kesehatan dan dukun. Fenomena dukun beranak menjadi fenomena yang menarik, karena nilai-nilai kearifan lokal yang tetap terjaga. Penelitian ini bertujuan mengetahui lebih dalam pengalaman partisipan melahirkan di rumah dengan kemitraan tenaga kesehatan dan dukun beranak. Metode yang digunakan dengan desain kualitatif dengan pendekatan fenomenologi. Penentuan partisipan menggunakan metode purposive sampling. Metode pengumpulan data dengan wawancara mendalam pada ibu dan FGD pada Kepala Desa, Pemangku Adat, dukun beranak dan tenaga kesehatan. Hasil penelitian ini diperoleh 6 tema yaitu pilihan tempat dan penolong saat melahirkan, keyakinan ibu terhadap dukun beranak dalam proses melahirkan, tradisi manyirira dan berurut selama hamil dan setelah melahirkan, perasaan ibu dengan kemitraan, tugas mandiri dukun beranak setelah melahirkan dan Pemanfaatan pelayanan kesehatan. Disarankan pada tenaga kesehatan untuk memiliki pengetahuan tentang budaya yang dipercayai oleh pasien dan keluarga, agar bisa memberikan dampak positif terhadap peningkatan kualitas pelayanan kesehatan ibu melahirkan dalam konteks budaya.

Keywords : Traditional birth attendants, Partnership, Experience of mother, Health workers.

Kata Kunci : Dukun beranak, Kemitraan, Pengalaman ibu, Tenaga Kesehatan.

Correspondence : Awang Saputra, Jln. Warung Jati Barat No. 17 Pancoran Kalibata Jakarta Selatan 12740
Email : awangsaputra1970@gmail.com , 0812 8231 950

• Received 24 November 2019 • Accepted 27 April 2020 • p - ISSN : 2088-7612 • e - ISSN : 2548-8538 •

DOI: <https://doi.org/10.25311/keskom.Vol6.Iss1.480>

INTRODUCTION

The efforts to preserve life by maintaining healthy, it must have a positive behavior where one of them about cervical cancer is not a disease that needs to be feared. The key to efforts to cure this type of cancer is early detection. Only 5% of women do early detection of cervical cancer with a Pap smear while ideally is 80% of the population of Indonesian women. In 2013 Banten Province, Wahidin (2013) was estimated the number of cervical cases 2,252 peoples, with 35 providers, 600 screening officers, and 6 trainers.

Culture of shame for inspection of reproductive organs, lower self-motivation early detection of cervical cancer, lower knowledge about cervical cancer and lower levels of education, cravings, and many indicators causes to do early detection of cervical cancer. Rasjidi (2009) said that primary prevention is an activity that can be done by everyone to avoid themselves from factors that can cause cancer. One part of primary prevention is to provide the human papillomavirus (HPV) vaccine, administering an HPV vaccine will eliminate HPV infection.

Early detection or screening is secondary prevention of cervical cancer to find early cases so that healing can be improved and to slow down or stop the disease at an early stage. Suwiyoga (2009) states this is done through early diagnosis of dysplasia in a variety of clinical and laboratory ways, one of which is a Papanicolaou smear or known as a Pap smear. Cervical cancer is cancer that occurs in the cervix of the female reproductive area which is the entrance to the uterus which is located between the uterus with the hole's intercourse (Diananda, 2012).

The benefits of Pap smear, Sumaryati (2003) said to detect early on the existence of inflammation of the uterus and the level of inflammation, the presence of degenerative abnormalities in the uterus, the presence/absence of signs of malignancy (cancer) in the uterus. Another statement Lestadi (2009) said the benefits of Pap smears for sito hormonal evaluation, diagnosing inflammation, identification of organisms that cause inflammation, diagnosing pre-cancerous disorders (dysplasia) of the early or advanced cervix (in situ/in fasive carcinoma).

Pap smear test is technically by taking a mucus sweep using a spatula or a kind of soft brush. Cervical mucus is taken by a doctor or midwife to be smeared and fixed (fixed) on a glass slide. Then by using a microscope, a cytologist will test the uterine cells (Nurhasanah, 2012).

Notoatmodjo (2008) states that knowledge is the result of "knowing" and this happens after people sensing a certain object. Sensing occurs through the human senses, which are the sense of sight, hearing, taste, and touch. Knowledge of cognition is a very important domain for the formation of one's actions (overt behavior). Motivation is an impetus found in a person to try to make changes in behavior that is better in meeting their

needs. Motivation occurs when someone has the desire and willingness to carry out an activity or action to achieve certain goals (Uno, 2012).

The marriages more than 20 years have a greater risk of changing cervical cells. Thus, the longer the age of marriage the higher the intensity of sexual intercourse this indicates the more vulnerable the risk to cervical cancer (Fitria, 2007).

The number of children is someone who has given birth to a baby that can live, where if the number of children is more than two people or the distance of labor is too close having a risk of changes in abnormal cells in the cervix. If the number of children causes abnormal cell changes from the epithelium in the cervix which can develop in malignancies.

Age is a unit of time that measures the time of existence of an object or creature, both living and dead, so that age is measured by the date it was born so that it dates. concerning this age where one is seen from an adult.

Adler (2011) said the education is the process by which all human abilities (talents and abilities acquired) that can be influenced by habituation, are enhanced by good habits through means that are artistically created and used by anyone to help others or themselves achieve the goals set that are habits the good.

Limitation of research is early detection of cervical cancer behavior with a Pap smear method on working women's as national health insurance participant as a dependent variable because there are too many women does not cervical cancer detection with a Pap smear method even though the cost of the examination is borne by the health insurance provider national and incidentally come from educational institutions in the health sector and work in the health sector.

The relationship of women's behavior in conducting a Pap smear inspection is the level of knowledge, women's attitudes, women's perceptions, family social support, attitudes of health workers, availability of service facilities, economic and time factors (Sari, 2012).

Pap smears can already find cervical cancer, although there is still a pre-cancerous level (early stage) so that it can provide 100% hope of recovery. As a tool for early diagnosis of cervical cancer has been carried out since the last three decades. In developed countries, it has been shown to reduce the incidence of invasive cervical cancer by 46-76% and mortality of cervical cancer by 50-60% (Tavassoli & Devilee, 2012).

Pap smear inspection is performed on women aged 25-50 years or married but for the most vulnerable to cervical cancer aged between 35-55 years, this is used because of the many conveniences that are affordable prices but specifically civil servants do not need to pay because it has guaranteed through health insurance, easy to diagnose and diagnose valid until the results.

This research is different from the others, the sample is based on a proportional random technique using the Slovin formula, respondents are women's work as civil servants in the Health Service, the background knowledge and education about health, the location is determined in Serang City, Banten Province as one of the parameters of a working woman aware for their reproductive health. Cross-sectional analysis techniques, with univariate, bivariate, and multivariate tests. The findings are 6 variables are related (knowledge, self-motivation, age, length of the marriage, number of children and education), on early detection of cervical cancer behavior with pap smear inspection.

The limitation of this study is the cross-sectional study design, the variables in this study are only based on the conceptual framework, the sample used is only limited to women's work in the Serang City Health Services, and the correctness of the data is highly dependent on the level of honesty of the respondents

METHOD

The method of this study is an analytic survey with a cross-sectional design, which aims to determine the relationship between independent and dependent variables at the same time (Notoatmodjo, 2010). The target population of women's work as female civil servant workers in the Serang City Health Office is 361 people. The affordable population of women's work as civil servants aged 30-55 years and had married 293 peoples The sampling technique uses non-probability sampling with purposive sampling, the technique of data collection uses, and documentation studies.

Univariate analysis data analysis techniques to determine the picture (in the form of numbers and percentages), including the behavior of cervical cancer early detection by the Pap smear method, knowledge, self-motivation, age, length of the marriage, number of children and education. Bivariate analysis to determine the relationship between variables with the chi-square test. Multivariate analysis to determine the relationship of independent variables and some of its compounding variables using logistic regression because the independent variables are of the categoric type.

RESULTS

Univariate analysis show that never cervical cancer detection 45.57%, and had detection 54.43%, never done a pap smear inspection 62.03%, and had inspection 37.97%, lower knowledge 3.17 %, and higher knowledge 96.83%, lower self-motivation 49.11%, and higher self-motivation 51.89%, aged 30-42 years 84.82% and 43-55 years 15.8%, length of marriage <10 years 48.10%, 11-20 years 42.40% and > 20 years 9.50%, don't have child of 25.32%, 1-2 children 47.46% and more than 3 children 27.22%, educated in high school & diploma 41.78% and 58.22%

undergraduate education. The results of univariate show at table 1:

Table 1 : Univariate Analysis

No	Variable	Category	n	%
1	Cervical Cancer Inspection Behavior	Ever	86	54,43
		Never	72	45,57
1a	Pap Smear Inspection Behavior	Ever	98	62,03
		Never	60	37,97
2	Knowledge	Lower	5	3,17
		Higher	153	96,83
3	Self-Motivation	Lower	76	49,11
		Higher	82	51,89
4	Age	30-42 Year	134	84,82
		43-55 Year	24	15,18
5	Length of Marriage	< 10 Year	76	48,10
		11-20 Year	67	42,40
		> 20 Year	15	9,50
6	Number of Children	Nothing	40	25,32
		1-2 persons	75	47,46
		< 3 persons	43	27,22
7	Education	SMA, Diploma	66	41,78
		Graduates	92	58,22

Bivariate analysis carried out aims to determine the relationship between the independent variables are knowledge, self-motivation, length of the marriage, age, number of children, and education with the dependent variable cervical cancer detection behavior with pap smear inspection. The results of bivariate show at table 2 :

Table 2 : Bivariate Analysis

Variable	Category	Pap Smear Inspection Behavior				Total		p - value	OR (Odd Ratio)
		Never		Ever		n	%		
		n	%	n	%				
Knowledge	Lower	14	8,9	15	9,5	29	18,4	0,046	0,642
	Higher	84	53,1	45	28,5	129	81,6		
Self-Motivation	Lower	41	25,9	38	24,1	79	50,0	0,004	0,933
	Higher	57	36,1	22	13,9	79	50,0		
Age	30-42 Year	89	56,3	45	28,5	134	84,8	0,007	3,296
	43-55 Year	9	5,7	15	9,5	24	15,2		
Length of Marriage	< 10 Year	64	40,5	11	7,0	75	47,5	0,000	4,294
	11-20 Year	28	17,7	39	24,7	67	40,4		
	> 20 Year	6	3,8	10	6,3	16	10,1		
Number of Children	Nothing	29	18,4	12	7,6	41	26,0	0,017	1,795
	1-2 persons	50	31,6	24	15,2	74	41,8		
	< 3 persons	19	12	24	15,2	43	27,2		
Education	SMA, Diploma	52	32,9	14	8,9	66	41,8	0,000	3,714
	Graduates	46	29,1	46	29,1	92	58,2		

Relationship of knowledge with cervical cancer detection behavior with pap smear inspection, 29 people with lower knowledge 8.9% never detection and 9.5%, had detection with a pap smear inspection, while 129 people with higher knowledge 53.1% never detection and 28.5% had detection with a pap smear inspection. Statistical test value of p=0.046 means that p<(α=0.05) concluded that there is a relationship between knowledge and cervical cancer detection behavior with pap smear inspection.

Relationship between self-motivation and cervical cancer detection behavior with pap smear inspection, 79 people with lower motivation 25.9% never detection, and 24.1% had detection, whereas 79 people with higher motivation 36.1% never detection and 13.9% had detection. Statistical test value of p=0.004 means that p<(α=0.05) concluded that there is a relationship between self-motivation and cervical cancer

detection behavior with pap smear inspection.

Relationship between age and cervical cancer detection behavior with pap smear inspection, 134 people aged 30-42 years 56.3% never detection and 28.5% had detection, while 24 people aged 43-55 years, 5.7% never detection and 9.5% had detection. The statistical value of $p = 0.007$ means that $p < (\alpha = 0.05)$, concluded that there is a relationship between age and cervical cancer detection behavior with pap smear inspection.

Relationship between long marriage and cervical cancer detection behavior with pap smear inspection, 75 people long married <10 years 40.5% never detection and 7.0% had detection, while 67 people were married long 11-20 years, 17.7% never detection and 24.7% had detection, then 16 people long married > 20 years 3.8% never detection and 6.3% detection. Statistical test obtained the value of $p = 0,000$ meaning $p < (\alpha = 0.05)$, concluded that there was a relationship between the length of the marriage and cervical cancer detection behavior with pap smear inspection.

The relationship the number of children with cervical cancer detection behavior with pap smear inspection, 41 people have not children 18.4% never detection and 7.6% had detection, while 74 people have birth to 1-2 children, 31.6% never detection and 15.2% had detection, then 43 more than 3 children, 12% never detection and 15.2% had detection. Statistical test obtained the value of $p = 0.017$ means that $p < (\alpha = 0.05)$ concluded that there is a relationship between the number of children with cervical cancer detection behavior with Pap smear inspection.

Relationship of education with cervical cancer detection behavior with pap smear inspection, 66 people with lower education 32.9% never detection, and 8.9% had detection, while 92 people with higher education 29.1% never detection, and 29.1% had detection. Statistical test obtained the value of $p = 0,000$ means that $p < (\alpha = 0.05)$ concluded that there is a relationship between education with cervical cancer detection behavior with pap smear inspection.

Multivariate analysis begins with the determination of candidate variables that will be included in multivariate analysis with p -values < 0.25, are knowledge (0.046), number of children (0.017), age (0.007), self-motivation (0.004), length of marriage (0,000), and education (0,000), then subsequently included in multivariate analysis together with a significant value limit $p < 0.05$, the selection of models is gradual using all the independent variables that have passed the sensor entered in the model starting from the p -value that is the biggest, it showed at table 3:

Table 3: Regression Logistic Analysis

Variable	p-value
Knowledge	0,046
Number of Children	0,017

Age	0,007
Self Motivation	0,004
Education	0,000
Length of Marriage	0,000

After going through several models, we can know the final results of multivariate analysis, which final modeling results of logistic regression analysis (multivariate), it turns out that all variables have a value of $p < 0.05$, so it can be concluded knowledge variables ($p = 0.046$; $OR = 0.642$), self-motivation ($p = 0.004$; $OR = 0.933$), age ($p = 0.007$; $OR = 3,296$), length of marriage ($p = 0,000$; $OR = 4,294$), number of children ($p = 0,017$; $OR = 1,795$), and education ($p = 0,000$; $OR = 3,714$) are significantly related to examination behavior pap smear, and the length of marriage variable is the most dominant variable ($p = 0,000$; $OR = 4,294$) related to cervical cancer detection behavior with pap smear inspection, meaning that working woman whose marriage age is longer will have the opportunity to conduct cervical cancer detection behavior with pap smear inspection, by 4.294 times compared to a working woman whose marriage age is not long ago, it showed at table 4:

Table 4: Regression Logistic Analysis

Variable	β	p-value	OR
Knowledge	3,090	0,046	0,642
Self Motivation	1,826	0,004	0,933
Age	-0,682	0,007	3,296
Length of Marriage	-1,478	0,000	4,294
Number of Children	-1,105	0,017	1,795
Education	-1,312	0,000	3,714

DISCUSSION

Most respondents had cervical cancer detection and never had a Pap smear inspection. This fact provides information that respondents who incidentally are female employees in the Health Service whose daily activities are inseparable from the health sector, in terms of reproductive health care (inspection of cervical cancer and pap smears) are still low, it could be due to limited places or hours service. Notoatmodjo (2007) said caused someone to behave or perform a certain action, among others, because there are thoughts and feelings in the form of knowledge, attitudes, perceptions, beliefs, and the presence of important figures as role models. Supported by the available resources including are facilities, costs, access, time, energy, and culture.

Pap smear inspection is a preventive measure to conduct early detection and to find out health problems, especially cervical cancer. Although there have been many studies conducted on cervical cancer, but only a small part can be overcome. This situation reflects that health problems that have emerged to the surface of human reach are only a small part and there will be many that have not been reached or addressed. Pap smear examination is also a form of improving healthy living

easier to develop themselves to make decisions and take action. The higher the education of a woman, the more potential it is to conduct a healthy behavior such as carrying out a Pap Smear inspection in detecting abnormalities in the reproductive area.

CONCLUSIONS

More women's work they had cervical cancer inspection, never early detection with a Pap smear, higher knowledge, higher self-motivation, age 30-42 years old, married for 11-20 years, had 1-2 children and higher education. There is a significant relationship between Pap smear inspection behavior with knowledge, self-motivation, age, length of the marriage, number of children, and education. The length of the marriage is the most dominant factor. As working women must be increased activation on early cervical cancer detection behavior with Pap smears inspection.

Acknowledgement

In writing this paper, many people supported me either directly or indirectly. Special thanks to Mr. Toyalis as head health service office of Serang city who provide policy support my research, Mrs. Suginarti and Mr. Ahmad Faridi as lectures at public health postgraduate program University Muhammadiyah of Prof.Dr.Hamka who provided technical guidance and direction in completing my research.

Author's Contributions

The author solely wrote this paper without help except language editing and polishing and also formatted of the final manuscript.

Ethics

I have correctly followed the expected research ethics in writing this paper.

REFERENCES

- Adler Mortimer, 2011. How to Read A Book: The Classic Guide To Intelligent Reading. A Touchstone Book, New York. ISBN : 0671212095, pp: 11.
- Ajeng Novita Sari, 2017. Hubungan Motivasi Deteksi Dini Kanker Serviks Dengan Tindakan Pap smear Pada Wanita Usia Subur di Wilayah Gonilan, Indonesian Journal on Medical Science 2017, 4(2): 189-195. <http://ejournal.ijmsbm.org/index.php/ijms/article/view/115/115>. (Accessed on December 17, 2018)
- Darnindro N, et al., 2006. Pengetahuan Sikap Perilaku Perempuan yang Sudah Menikah Mengenai Pap smear dan Faktor-Faktor yang Berhubungan di Rumah Susun Klender Jakarta. Jakarta: Fakultas Kedokteran Universitas Indonesia, Majalah Kedokteran Indonesia 2006, 57(7): 1-7. [https://docplayer.info/29682824-Pengetahuan-](https://docplayer.info/29682824-Pengetahuan-sikap-perilaku-perempuan-yang-sudah-menikah-mengenai-pap-smear-dan-faktor-faktor-yang-berhubungan-di-rumah-susun-klender-jakarta-2006.html)
- [sikap-perilaku-perempuan-yang-sudah-menikah-mengenai-pap-smear-dan-faktor-faktor-yang-berhubungan-di-rumah-susun-klender-jakarta-2006.html](https://docplayer.info/29682824-Pengetahuan-sikap-perilaku-perempuan-yang-sudah-menikah-mengenai-pap-smear-dan-faktor-faktor-yang-berhubungan-di-rumah-susun-klender-jakarta-2006.html). (Accessed on September 6, 2018)
- Diananda R, 2009. Kanker Serviks: Sebuah Peringatan Buat Wanita. In: Diananda, R. Mengenal Seluk-Beluk Kanker. Katahari, Yogyakarta. ISBN : 9792544437, pp: 43-60.
- Kristiana E dan Niniek S, 2017. Pengetahuan Sikap dan Praktik Wanita Usia Subur Dalam Pemeriksaan Pap smear di Desa Bangsri Kecamatan Kertosono Kabupaten Nganjuk Tahun 2015. Jurnal Ilmiah Bidan 2017, 2(1) : 18 - 24 . <https://ejournal.ibi.or.id/index.php/jib/article/view/21/19>. (Accessed on September 6, 2018)
- Erma R, 2018. Hubungan Pengetahuan dan Sikap Dengan Pemeriksaan Pap smear Pada Pasangan Usia Subur (PUS) di Desa Joho Kecamatan Wates Kabupaten Kediri, Journal for Quality in Women's Health 2018, 1(1) : 10 - 15 . <https://media.neliti.com/media/publications/268128-hubungan-pengetahuan-dan-sikap-dengan-pe-51679d9d.pdf>. (Accessed on September 6, 2018)
- Fitria A, 2007. Panduan Lengkap Kesehatan Wanita. Gala Ilmu, Yogyakarta. ISBN : 979259793x, pp: 79.
- Hakimah U, 2016. Hubungan Usia Menikah dan Paritas Dengan Tindakan Pap-Smear Di Yayasan Kanker, Wisnuwardhana: Jurnal Berkala Epidemiologi 2016, 4(3) : 420 - 431 . <https://media.neliti.com/media/publications/94929-ID-none.pdf>. (Accessed on September 6, 2018)
- Hasbullah, 2011. Dasar-Dasar Ilmu Pendidikan. Rajawali Press, Jakarta. ISBN : 9794216933, pp: 78
- Imam Rasjidi, 2009. Epidemiologi Kanker Serviks. Indonesian Journal of Cancer 2009, 3(3); 103-109. . (Accessed on September 3, 2018)
- Lestadi L, 2009. Sitologi Pap Smear. EGC, Jakarta. ISBN : 9794489963, pp: 79
- Notoatmodjo Soekidjo, 2014. Ilmu Perilaku Kesehatan. Rineka Cipta, Jakarta. ISBN : 9789795189831, pp: 45
- Notoatmodjo Soekidjo, 2010. Metode Penelitian Kesehatan. Rineka Cipta, Jakarta. ISBN : 9789795189848, pp: 89
- Notoatmodjo Soekidjo, 2007. Pendidikan Kesehatan dan Ilmu Perilaku. Rineka Cipta, Jakarta. ISBN : 9789790980327, pp: 98
- Nurhasanah Cut, 2008. Pengaruh Karakteristik dan Perilaku Pasangan Usia Subur (PUS) Terhadap Pemeriksaan Pap Smear di Banda Aceh Tahun 2008. Sekolah Pascasarjana Universitas Sumatra Utara. 2008: 103.

- <http://repository.usu.ac.id/bitstream/handle/123456789/6762/08E00456.pdf?sequence=1&isAllowed=y>. (Accessed on September 3, 2018)
- Reni H, 2018. Hubungan Pengetahuan Wanita Terhadap Pemeriksaan Papsmear di Wilayah Puskesmas Garuda Kota Pekanbaru. *Jurnal Endurance* 2018, 3 (3) : 5 9 6 - 6 0 2 . DOI : <http://doi.org/10.22216/jen.v3i3.3451>
- Sari Ratna Puspita Arum, 2008. Faktor-Faktor Yang Berhubungan Dengan Perilaku Wanita Dalam Melakukan Pemeriksaan Pap Smear di Kelurahan Tembalang Semarang. Program Studi Ilmu Keperawatan Fakultas Kedokteran Universitas Diponegoro Semarang. 2008: 1. <http://eprints.undip.ac.id/16070/1/Ratna.pdf>. (Accessed on September 23, 2018)
- Sri Widiani NK, 2014. Hubungan Motivasi Dengan Tindakan Pap Smear Pada Wanita Usia Subur di Wilayah Kerja Puskesmas II Denpasar Barat, *Jurnal Ilmu Keperawatan FK Universitas Udaya Bali* 2014, 2(3): 1-7. <https://ojs.unud.ac.id/index.php/coping/article/view/10773>. (Accessed on September 13, 2018)
- Sumaryati Aryoso, 2003. Pengembangan Pola Operasional Dalam Percepatan Penurunan Angka Kematian Ibu dan Anak, Program Litbang Dalam Mendukung Percepatan Penurunan Angka Kematian Ibu dan Anak, Policy paper. Badan Penelitian dan Pengembangan Kesehatan, Departemen Kesehatan: Jakarta. http://repository.litbang.kemkes.go.id/3012/1/20-99Z_Book%20Manuscript-36-1-10-2.pdf. (Accessed on September 3, 2018)
- Sunaryo, 2013. Psikologi Untuk Keperawatan. EGC, Jakarta. ISBN : 9789790443631, pp. 98
- Suwiyoga K, 2007. Kanker Serviks: Penyakit Keganasan Fatal yang dapat di Cegah, *Majalah Obstetri dan Ginekologi Indonesia* 2007, 31(1): 3-25. <http://inajog.com/index.php/journal/article/view/112/107>. (Accessed on November 3, 2018)
- Tavassoli FA, & Devilee P, 2003. Pathology & Genetics of Tumors of The Breast and Female Genital Organ. IARC Press, Lyon. ISBN : 9789283224129, pp: 259-271.
- Ulfiana E, 2013. Analisis Faktor yang Memengaruhi Niat Wanita Pasangan Usia Subur untuk Pap smear di Wilayah Kelurahan Kedung Mundu Wilayah Puskesmas Kedungmundu Kota Semarang, *Jurnal Kebidanan* 2013, 2(4): 17-23. DOI: <http://Dx.Doi.Org/10.31983/Jkb.V2i4.100>
- Uno Hamzah B, 2007. Teori Motivasi dan Pengukurannya: Analisis di Bidang Pendidikan. Bumi Aksara, Jakarta. ISBN : 9789790100046, pp: 9
- Wahidin Mugi, 2013. Deteksi Dini Kanker Rahim dan Payudara di Indonesia 2007-2013. *Buletin Jangkauan Data dan Informasi Kemenkes RI*: Jakarta, pp: 78.